

Oct. 03 2005: Lecture 8:

Complex Numbers and Euler's Formula

Reading:

Kreyszig Sections: §12.1 (pp:652–56), §12.2 (pp:657–62), §12.6 (pp:679–82), §12.7 (pp:682–85)

Complex Numbers and Operations in the Complex Plane

With $i \equiv \sqrt{-1}$, the complex numbers can be defined as the space of numbers spanned by the vectors:

$$\begin{pmatrix} 1 \\ 0 \end{pmatrix} \text{ and } \begin{pmatrix} 0 \\ i \end{pmatrix} \quad (8-1)$$

so that any complex number can be written as

$$z = x \begin{pmatrix} 1 \\ 0 \end{pmatrix} + y \begin{pmatrix} 0 \\ i \end{pmatrix} \quad (8-2)$$

or just simply as

$$z = x + iy \quad (8-3)$$

where x and y are real numbers. $\operatorname{Re} z \equiv x$ and $\operatorname{Im} z \equiv y$.

MATHEMATICA® Example: (notebook) Lecture-08
Operations on complex numbers

Addition, subtraction, multiplication, division

@@ Complex Plane and Complex Conjugates

Because the complex basis can be written in terms of the vectors in Equation 8-1, it is natural to plot complex numbers in two dimensions—typically these two dimensions are the “complex plane” with $(0, i)$ associated with the y -axis and $(1, 0)$ associated with the x -axis.

The reflection of a complex number across the real axis is a useful operation. The image of a reflection across the real axis has some useful qualities and is given a special name—“the complex conjugate.”




Figure 8-1: Plotting the complex number z in the complex plane: The complex conjugate (\bar{z}) is a *reflection* across the *real axis*; the minus $(-z)$ operation is an *inversion* through the origin; therefore $-(\bar{z}) = (-z)$ is equivalent to either a reflection across the *imaginary axis* or an *inversion* followed by a reflection across the *real axis*.

The real part of a complex number is the projection of the displacement in the *real direction* and also the average of the complex number and its conjugate: $\text{Re } z = (z + \bar{z})/2$. The imaginary part is the displacement projected onto the *imaginary axis*, or the complex average of the complex number and its reflection across the *imaginary axis*: $\text{Im } z = (z - \bar{z})/(2i)$.

1. Polar Form of Complex Numbers

There are physical situations in which a transformation from Cartesian (x, y) coordinates to polar (*or cylindrical*) coordinates (r, θ) simplifies the algebra that is used to describe the physical problem.

An equivalent coordinate transformation for complex numbers, $z = x+iy$, has an analogous simplifying effect for *multiplicative operations* on complex numbers. It has been demonstrated how the complex conjugate, \bar{z} , is related to a reflection—multiplication is related to a **counter-clockwise** rotation in the complex plane. Counter-clockwise rotation corresponds to increasing θ .

The transformations are:

$$(x, y) \rightarrow (r, \theta) \begin{cases} x = r \cos \theta \\ y = r \sin \theta \end{cases} \quad (8-4)$$

$$(r, \theta) \rightarrow (x, y) \begin{cases} r = \sqrt{x^2 + y^2} \\ \theta = \arctan \frac{y}{x} \end{cases}$$

where $\arctan \in (-\pi, \pi]$.

⊗ Multiplication, Division, and Roots in Polar Form

One advantage of the polar complex form is the simplicity of multiplication operations:

DeMoivre's formula:

$$z^n = r^n(\cos n\theta + i \sin n\theta) \quad (8-5)$$

$$\sqrt[n]{z} = \sqrt[n]{z} \left(\cos \frac{\theta + 2k\pi}{n} + i \sin \frac{\theta + 2k\pi}{n} \right) \quad (8-6)$$

MATHEMATICA® Example: (notebook) Lecture-07

Polar Form of Complex Numbers

Writing a function to convert to polar form

Exponentiation and Relations to Trigonometric Functions

Exponentiation of a complex number is defined by:

$$e^z = e^{x+iy} = e^x(\cos y + i \sin y) \quad (8-7)$$

Exponentiation of a purely imaginary number advances the angle by rotation:

$$e^{iy} = \cos y + i \sin y \quad (8-8)$$

combining Eq. 8-8 with Eq. 8-7 gives the particularly useful form:

$$z = x + iy = re^{i\theta} \quad (8-9)$$

and the useful relations (that can be obtained simply by considering the geometry of the complex plane)

$$e^{2\pi i} = 1 \quad e^{\pi i} = -1 \quad e^{-\pi i} = -1 \quad e^{\frac{\pi}{2}i} = i \quad e^{-\frac{\pi}{2}i} = -i \quad (8-10)$$

Judicious subtraction of powers in Eq. 8-8 and generalization gives the following useful relations for trigonometric functions:

$$\begin{aligned}
 \cos z &= \frac{e^{iz} + e^{-iz}}{2} & \sin z &= \frac{e^{iz} - e^{-iz}}{2i} \\
 \cosh z &= \frac{e^z + e^{-z}}{2} & \sinh z &= \frac{e^z - e^{-z}}{2} \\
 \cos iz &= \cosh z & i \sin z &= \sinh iz \\
 \cos iz &= \cosh z & \sin iz &= i \sinh z
 \end{aligned} \tag{8-11}$$

MATHEMATICA® Example: (notebook) Lecture-07

Numerical precision and rounding of complex numbers

Numerical and symbolic representations of complex numbers

Roots of polynomial equations

Handling complex roots of polynomial equations