

Dec. 07 2005: **Lecture 27:****Eigenfunction Basis**

Reading:

Kreyszig Sections: §4.7 (pp:233–38) , §4.8 (pp:240–248)

**Sturm-Liouville Theory, Orthogonal Eigenfunctions**

The trigonometric functions have the property that they are orthogonal, that is:

$$\begin{aligned} \int_{x_0}^{x_0+\lambda} \sin\left(\frac{2\pi M}{\lambda}x\right) \sin\left(\frac{2\pi N}{\lambda}x\right) dx &= \begin{cases} \frac{\lambda}{2} & \text{if } M = N \\ 0 & \text{if } M \neq N \end{cases} \\ \int_{x_0}^{x_0+\lambda} \cos\left(\frac{2\pi M}{\lambda}x\right) \cos\left(\frac{2\pi N}{\lambda}x\right) dx &= \begin{cases} \frac{\lambda}{2} & \text{if } M = N \\ 0 & \text{if } M \neq N \end{cases} \\ \int_{x_0}^{x_0+\lambda} \cos\left(\frac{2\pi M}{\lambda}x\right) \sin\left(\frac{2\pi N}{\lambda}x\right) dx &= 0 \text{ for any integers } M, N \end{aligned} \quad (27-1)$$

This property allowed the Fourier series to be obtained by multiplying a function by one of the basis functions and then integrating over the domain.

 extra notes: *Inner Products for Functions* .....

The orthogonality relation for the trigonometric functions requires two things:

**Range** The range over which the functions are defined (i.e., values of  $x$  for which  $f(x)$  and  $g(x)$  have a inner product defined) and integrated in their inner product definition.

**Inner product** The projection operation of one function onto another.

For the trigonometric functions, the inner product was a fairly obvious choice:

$$f(x) \cdot g(x) = \int_0^{2\pi} f(x)g(x)dx \quad (27-2)$$

This inner product follows from the *l2-norm* for functions:

$$|f(x)| = \sqrt{\int f(x)f(x)dx} = \sqrt{\int f^2 dx} \quad (27-3)$$

which is one of the obvious ways to measure “the distance of a function from zero.” The l2-norm is employed in least-squares-fits.

However, there are different choices of inner products. For example, the Laguerre polynomials (or Laguerre functions)  $L_n(x)$  defined by

$$L_n(x) = \frac{e^x}{n!} \frac{d^n}{dx^n} (x^n e^{-x}) \quad (27-4)$$

for  $0 < x < \infty$  have the orthogonality relation for a weighted inner product:

$$L_n(x) \cdot L_m(x) = \int_0^\infty e^{-x} L_n(x) L_m(x) dx = \delta_{mn} \quad (27-5)$$

There are many other kinds of functional norms.

Many ordinary differential equations—including the harmonic oscillator, Bessel, and Legendre—can be written in a general form:

$$\frac{d}{dx} \left[ r(x) \frac{dy}{dx} \right] + [q(x) + \lambda p(x)] y(x) = 0 \quad (27-6)$$

which is called the *Sturm-Liouville problem*. Solutions to this equation are called eigensolutions for an eigenvalue  $\lambda$ . The function  $p(x)$  appears in the orthonormality relation:

$$\int p(x) y_{\lambda_1}(x) y_{\lambda_2}(x) dx = 0 \quad \text{if } \lambda_1 \neq \lambda_2 \quad (27-7)$$

The same "trick" of multiplying a function by one of the eigensolutions and then summing a series can be used to generate series solutions as a superposition of eigensolutions.

MATHEMATICA® Example: (notebook) Lecture-27

**Legendre functions**

*Expanding a function as a series of Legendre functions*

---



---



---



---



---