

Sept. 27 2006

Lecture 6: Linear Algebra I

Reading:

Kreyszig Sections: 7.5, 7.6, 7.7, 7.8, 7.9 (pages 302–305, 306–307, 308–314, 315–323, 323–329)

Vectors

Vectors as a list of associated information

$$\vec{x} = \begin{pmatrix} \text{number of steps to the east} \\ \text{number of steps to the north} \\ \text{number steps up vertical ladder} \end{pmatrix} \quad (6-1)$$

$$\vec{x} = \begin{pmatrix} 3 \\ 2.4 \\ 1.5 \end{pmatrix} \quad \text{determines position} \quad \begin{pmatrix} x_{\text{east}} \\ x_{\text{north}} \\ x_{\text{up}} \end{pmatrix} \quad (6-2)$$

The vector above is just one example of a position vector. We could also use coordinate systems that differ from the Cartesian (x, y, z) to represent the location. For example, the location in *cylindrical coordinate system* could be written as

$$\vec{x} = \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} r \cos \theta \\ r \sin \theta \\ z \end{pmatrix} \quad (6-3)$$

as a *Cartesian vector* in terms of the cylindrical coordinates (r, θ, z) .

The position could also be written as a cylindrical, or *polar* vector

$$\vec{x} = \begin{pmatrix} r \\ \theta \\ z \end{pmatrix} = \begin{pmatrix} \sqrt{x^2 + y^2} \\ \tan^{-1} \frac{y}{x} \\ z \end{pmatrix} \quad (6-4)$$

where the last term is the polar vector in terms of the Cartesian coordinates. Similar rules would apply for other coordinate systems like spherical, elliptic, etc.

However, vectors need not represent position at all, for example:

$$\vec{n} = \begin{pmatrix} \text{number of Hydrogen atoms} \\ \text{number of Helium atoms} \\ \text{number of Lithium atoms} \\ \vdots \\ \text{number of Plutonium atoms} \\ \vdots \end{pmatrix} \quad (6-5)$$

Scalar multiplication

$$\frac{1}{N_{\text{avag.}}} \vec{n} \equiv \begin{pmatrix} \frac{\text{number of H}}{N_{\text{avag.}}} \\ \frac{\text{number of He}}{N_{\text{avag.}}} \\ \frac{\text{number of Li}}{N_{\text{avag.}}} \\ \vdots \\ \frac{\text{number of Pu}}{N_{\text{avag.}}} \\ \vdots \end{pmatrix} = \begin{pmatrix} \text{moles of H} \\ \text{moles of He} \\ \text{moles of Li} \\ \vdots \\ \text{moles of Pu} \\ \vdots \end{pmatrix} = \vec{m} \quad (6-6)$$

Vector norms

$$\|\vec{x}\| \equiv x_1^2 + x_2^2 + \dots x_k^2 = \text{euclidean separation} \quad (6-7)$$

$$\|\vec{n}\| \equiv n_{\text{H}} + n_{\text{He}} + \dots n_{132?} = \text{total number of atoms} \quad (6-8)$$

Unit vectors

unit direction vector

$$\hat{x} = \frac{\vec{x}}{\|\vec{x}\|}$$

mole fraction composition

$$\hat{m} = \frac{\vec{m}}{\|\vec{m}\|}$$

(6-9)

(6-10)

Extra Information and Notes

Potentially interesting but currently unnecessary

If $\mathbb{R}$ stands for the set of all real numbers (i.e., 0, -1.6 , $\pi/2$, etc.), then can use a shorthand to specify the position vector, $\vec{x} \in \mathbb{R}^N$ (e.g., each of the N entries in the vector of length N must be a real number—or in the set of real numbers. $\|\vec{x}\| \in \mathbb{R}$).

For the unit (direction) vector: $\hat{x} = \{\vec{x} \in \mathbb{R}^3 \mid \|\vec{x}\| = 1\}$ (i.e., the unit direction vector is the set of all position vectors such that their length is unity—or, the unit direction vector is the subset of all position vectors that lie on the unit sphere. $\vec{x}$ and $\hat{x}$ have the same number of entries, but compared to $\vec{x}$, the number of independent entries in $\hat{x}$ is smaller by one.

For the case of the composition vector, it is strange to consider the case of a negative number of atoms, so the mole fraction vector $\vec{n} \in (\mathbb{R}^+)^{\text{elements}}$ ($\mathbb{R}^+$ is the real non-negative numbers) and $\hat{m} \in (\mathbb{R}^+)^{(\text{elements}-1)}$.

Matrices and Matrix Operations

Consider methane (CH_4), propane (C_3H_8), and butane (C_4H_{10}).

$$\underline{M}_{HC} = \begin{pmatrix} \text{H-column} & \text{C-column} \\ \frac{\text{number of H}}{\text{methane molecule}} & \frac{\text{number of C}}{\text{methane molecule}} \\ \frac{\text{number of H}}{\text{propane molecule}} & \frac{\text{number of C}}{\text{propane molecule}} \\ \frac{\text{number of H}}{\text{butane molecule}} & \frac{\text{number of C}}{\text{butane molecule}} \end{pmatrix} \begin{matrix} \text{methane row} \\ \text{propane row} \\ \text{butane row} \end{matrix} \quad (6-11)$$

$$\underline{M_{HC}} = \begin{pmatrix} 4 & 1 \\ 8 & 3 \\ 10 & 4 \end{pmatrix} = \begin{pmatrix} M_{11} & M_{12} \\ M_{21} & M_{22} \\ M_{31} & M_{32} \end{pmatrix} \quad (6-12)$$

Matrices as a linear transformation of a vector

$$\vec{N_{HC}} = (\text{number of methanes, number of propanes, number of butanes}) \quad (6-13)$$

$$= (N_{HC\ m}, N_{HC\ p}, N_{HC\ b}) \quad (6-14)$$

$$= (N_{HC\ 1}, N_{HC\ 2}, N_{HC\ 3}) \quad (6-15)$$

$$(6-16)$$

$$N_{HC} \vec{M_{HC}} \equiv \sum_{i=1}^3 N_{HC\ i} M_{HC\ ij} = \vec{N} \quad (6-17)$$

The “summation” convention is often used, where a repeated index is summed over all its possible values:

$$\sum_{i=1}^p N_{HC\ i} M_{HC\ ij} \equiv N_{HC\ i} M_{HC\ ij} = N_j \quad (6-18)$$

For example, suppose

$$\vec{N}_{HC} = (1.2 \times 10^{12} \text{ molecules methane}, 2.3 \times 10^{13} \text{ molecules propane}, 3.4 \times 10^{14} \text{ molecules butane}) \quad (6-19)$$

$$\begin{aligned} \vec{N}_{HC} \underline{M}_{HC} &= \\ (1.2 \times 10^{14} \text{ methanes}, 2.3 \times 10^{13} \text{ propanes}, 3.4 \times 10^{12} \text{ butanes}) &\begin{pmatrix} \frac{4 \text{ atoms H}}{\text{methane}} & \frac{1 \text{ atoms C}}{\text{methane}} \\ \frac{8 \text{ atoms H}}{\text{propane}} & \frac{3 \text{ atoms C}}{\text{propane}} \\ \frac{10 \text{ atoms H}}{\text{butane}} & \frac{\text{atoms C}}{\text{butane}} \end{pmatrix} \\ &= (7.0 \times 10^{14} \text{ atoms H}, 2.0 \times 10^{14} \text{ atoms C}) \end{aligned} \quad (6-20)$$

Matrix transpose operations

Above the lists (or vectors) of atoms were stored as rows, often it is convenient to store them as columns. The operation to take a row to a column (and vice-versa) is a “transpose”.

$$\underline{M}_{HC}^T = \begin{pmatrix} \frac{\text{methane-column}}{\text{number of H}} & \frac{\text{propane-column}}{\text{number of H}} & \frac{\text{butane-column}}{\text{number of H}} \\ \frac{\text{methane molecule}}{\text{number of C}} & \frac{\text{propane molecule}}{\text{number of C}} & \frac{\text{butane molecule}}{\text{number of C}} \end{pmatrix} \begin{matrix} \text{hydrogen row} \\ \text{carbon row} \end{matrix} \quad (6-21)$$

$$\vec{N}_{HC}^T = \begin{pmatrix} \text{number of methanes} \\ \text{number of propanes} \\ \text{number of butanes} \end{pmatrix} = \begin{pmatrix} N_{HC\ m} \\ N_{HC\ p} \\ N_{HC\ b} \end{pmatrix} \quad (6-22)$$

$$\underline{M}_{HC}^T \vec{N}_{HC}^T = \vec{N}^T \begin{pmatrix} 4 & 8 & 10 \\ 1 & 3 & 4 \end{pmatrix} \begin{pmatrix} \text{number of methanes} \\ \text{number of propanes} \\ \text{number of butanes} \end{pmatrix} = \begin{pmatrix} \text{number of H-atoms} \\ \text{number of C-atoms} \end{pmatrix} \quad (6-23)$$

Matrix Multiplication

The next example supposes that some process produces hydrocarbons and be modeled with the pressure P and temperature T . Suppose (this is an artificial example) that the number of hydrocarbons produced in one millisecond can be related linearly to the pressure and temperature:

$$\begin{aligned}\text{number of methanes} &= \alpha P + \beta T \\ \text{number of propanes} &= \gamma P + \delta T \\ \text{number of butanes} &= \epsilon P + \phi T\end{aligned}\tag{6-24}$$

or

$$N_{HC}^T = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \\ \epsilon & \phi \end{pmatrix} \begin{pmatrix} P \\ T \end{pmatrix}\tag{6-25}$$

Lecture 06 MATHEMATICA® Example 1

Matrices

Download notebooks, pdfs, or html from <http://pruffle.mit.edu/3.016-2006>.

Here is an example operation that takes us from the processing vector $(P, T)^T$ to the number of hydrogens and carbons.

- 1: The matrix (Eq. 6-12) is entered as a list of sublists. The sub-lists are the rows of the matrix. The first elements of each row-sublist form the first column; the second elements are the second column and so on.

The **Length** of a matrix-object gives the number of row, and the second member of the result of **Dimensions** gives the number of columns.

All sublists of a matrix must have the same dimensions.

It is good practice to enter a matrix and then display it separately using **MatrixForm**. Otherwise, there is a risk of defining a symbol as a **MatrixForm**-object *and not as a matrix which was probably the intent*.

- 3: This command will generate an error.

Matrix multiplication in MATHEMATICA® is produced by the "dot" . operator. For matrix multiplication, $\underline{A}.\underline{B}$, the number of columns of $\underline{A}$ must be equal to the number of rows of $\underline{B}$.

- 5: The **Transpose** "flips" a matrix by producing a new matrix which has the original's i^{th} row as the new matrix's i^{th} column (or, equivalently the j^{th} column as the new j^{th} row). In this example, a 3×2 -matrix ($\underline{PTmatrix}$) is being left-multiplied by a 2×3 -matrix.

The resulting matrix would map a vector with values P and T to a vector for the rate of production of C and H.

```

1 MHC = {
  {4, 1},
  {8, 3},
  {10, 4}
}
MHC // MatrixForm

2 PTmatrix = {
  {α, β},
  {γ, δ},
  {ε, φ}
};
PTmatrix // MatrixForm

3 MPT = MHC. PTmatrix

4 Clear[MPT]

5 MPT = Transpose[MHC]. PTmatrix;
MPT // MatrixForm

```

Matrix multiplication is defined by:

$$\underline{AB} = \sum_i A_{ki} B_{ij}\tag{6-26}$$

The indices of the matrix defined by the multiplication $\underline{AB} = \underline{C}$ are C_{kj} .

Matrix Inversion


Sometimes what we wish to know, “What vector is it ($\vec{x}$), when transformed by some matrix ($\underline{A}$) gives us a particular result ($\vec{b} = \underline{A}\vec{x}$)?”

$$\begin{aligned}\underline{A}\vec{x} &= \vec{b} \\ \underline{A}^{-1}\underline{A}\vec{x} &= \underline{A}^{-1}\vec{b} \\ \vec{x} &= \underline{A}^{-1}\vec{b}\end{aligned}\tag{6-27}$$

The inverse of a matrix is defined as something that when multiplied with the matrix leaves a product that has no effect on any vector. This special product matrix is called the *identity matrix*.

$$\underline{A}\vec{x} = \vec{b}$$

$$\begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} b_1 \\ b_2 \end{pmatrix} \quad (6-28)$$



Lecture 06 MATHEMATICA® Example 3

Eliminating redundant equations or variables

Download notebooks, pdfs, or html from <http://pruffle.mit.edu/3.016-2006>.

Consider liquid water near the freezing point—dipole interactions will tend to make water molecules form clusters such as H_2O and H_4O_2 .

Then the mapping from molecules to the number of atoms becomes:

$$\begin{pmatrix} 2 & 4 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} N_{\text{H}_2\text{O}} \\ N_{\text{H}_4\text{O}_2} \end{pmatrix} = \begin{pmatrix} N_{\text{H}} \\ N_{\text{O}} \end{pmatrix} \quad (6-29)$$

This example treats this case where the columns are not linearly independent.

- 5: This equation is the same as the first row of $\underline{A}\vec{x}$ being set to the first entry of $\vec{b}$ for $\underline{A}\vec{x} = \vec{b}$.
- 7: This is an attempt to find the number of H_2O - and H_4O_2 -molecules, given the number of H- and O-atoms. Of course, it has to fail.
- 9: **Eliminate** produces a logical equality for each redundancy in a set of equations. In this case, the result expresses the fact that $2 \times$ (second row) is the same as the (first row).
- 10: The rank of a matrix, obtained with **MatrixRank**, gives the number of linearly independent rows.
- 11: The null space of a matrix, $\underline{A}$, is a linearly independent set of vectors $\vec{x}$, such that $\underline{A}\vec{x}$ is the zero-vector; this list can be obtained with **NullSpace**. The *nullity* is the number of vectors in a matrix's null space. The rank and the nullity must add up to the number of columns of $\underline{A}$

```

1  watmat = {{2, 4}, {1, 2}};
   watmat // MatrixForm
2  molvec = {h20, h402}
3  atomvec = {h, o}
4  atomvec // MatrixForm
5  eq[1] = (watmat.molvec)[[1]] == atomvec[[1]]
6  eq[2] = (watmat.molvec)[[2]] == atomvec[[2]]
7  Solve[{eq[1], eq[2]}, molvec]
8  ? Eliminate
9  Eliminate[{eq[1], eq[2]}, molvec]
10 MatrixRank[watmat]
11 NullSpace[watmat]
   Length[NullSpace[watmat]]

```