

Oct. 2 2006

Lecture 8: Complex Numbers and Euler's Formula

Reading:

Kreyszig Sections: 8.1, 8.2, 8.3 (pages 334–338, 340–343, 345–348)

Complex Numbers and Operations in the Complex Plane

Consider, the number zero: it could be *operationally* defined as the number, which when multiplied by any other number always yields itself; and its other properties would follow.

Negative numbers could be defined operationally as something that gives rise to simple patterns. Multiplying by -1 gives rise to the pattern $1, -1, 1, -1, \dots$ In the same vein, a number, \imath , can be created that doubles the period of the previous example: multiplying by \imath gives the pattern: $1, \imath, -1, -\imath, 1, \imath, -1, -\imath, \dots$ Combining the imaginary number, \imath , with the real numbers, arbitrarily long periods can be defined by multiplication; applications to periodic phenomena is probably where complex numbers have their greatest utility in science and engineering

With $i \equiv \sqrt{-1}$, the complex numbers can be defined as the space of numbers spanned by the vectors:

$$\begin{pmatrix} 1 \\ 0 \end{pmatrix} \text{ and } \begin{pmatrix} 0 \\ i \end{pmatrix} \quad (8-1)$$


so that any complex number can be written as

$$z = x \begin{pmatrix} 1 \\ 0 \end{pmatrix} + y \begin{pmatrix} 0 \\ i \end{pmatrix} \quad (8-2)$$

or just simply as

$$z = x + iy \quad (8-3)$$

where x and y are real numbers. $\operatorname{Re} z \equiv x$ and $\operatorname{Im} z \equiv y$.



Lecture 08 MATHEMATICA® Example 1

Operations on complex numbers

Download notebooks, pdfs, or html from <http://pruffle.mit.edu/3.016-2006>.

Straightforward examples of addition, subtraction, multiplication, and division of complex numbers are demonstrated. An example that demonstrates that MATHEMATICA® doesn't make *a priori* assumptions about whether a symbol is real or complex. An example function that converts a complex number to its polar form is constructed.

- 2: Just like `Pi` is a *mathematical constant*, the imaginary number is defined in MATHEMATICA® as something with the properties of i
- 3: Here, two numbers that are *potentially, but not necessarily* complex are defined.
- 4: Addition and multiplication are defined as for any symbol; here the results do not appear to be very interesting *because* the other symbols could themselves be complex...
- 5: And, `Simplify` doesn't help much even with assumptions.
- 6: The real and imaginary parts of a complex entity can be extracted with `Re` and `Im`. This demonstrates that MATHEMATICA® hasn't made assumptions about `a`, `b`, `c`, and `d`.
- 8-12 However, `ComplexExpand` does make assumptions that symbols are real and, here, demonstrate the rules for addition, multiplication, division, and exponentiation.
- 13: `Abs` calculates the magnitude (also known as modulus or absolute value) and `Arg` calculates the argument (or angle) of a complex number. Here, they are used to define a function to convert an expression to an equivalent *polar form of a complex number*.

```

1 imaginary = Sqrt[-1]
2 (-imaginary)^2
3 z1 = a + i b;
z2 = c + i d;
4 compadd = z1 + z2;
5 compmult = z1*z2;
6 Simplify[compmult,
a ∈ Reals && b ∈ Reals && c ∈ Reals && d ∈ Reals]
Mathematica doesn't assume that symbols are necessarily real...
7 Re[compadd]
Im[compadd]
However, the Mathematica function ComplexExpand does
assume that the variables are real...
8 ComplexExpand[Re[compadd]]
9 ComplexExpand[Im[compadd]]
10 ComplexExpand[Re[z1/z2]]
11 ComplexExpand[compmult]
12 ComplexExpand[Re[z1^3]]
ComplexExpand[Im[z1^3]]
Function to convert to Polar Form
13 Pform[z_] := Abs[z] Exp[i Arg[z]]
14 Pform[z1]
15 Pform[z1 /. {a → 2, b → -π}]
16 ComplexExpand[Pform[z1]]

```

Complex Plane and Complex Conjugates

Because the complex basis can be written in terms of the vectors in Equation 8-1, it is natural to plot complex numbers in two dimensions—typically these two dimensions are the “complex plane” with $(0, i)$ associated with the y -axis and $(1, 0)$ associated with the x -axis.

The reflection of a complex number across the real axis is a useful operation. The image of a

reflection across the real axis has some useful qualities and is given a special name—"the complex conjugate."




Figure 8-3: Plotting the complex number z in the complex plane: The complex conjugate (\bar{z}) is a *reflection* across the real axis; the minus $(-z)$ operation is an *inversion* through the origin; therefore $-(\bar{z}) = (\bar{-z})$ is equivalent to either a reflection across the imaginary axis or an inversion followed by a reflection across the real axis.

The real part of a complex number is the projection of the displacement in the real direction and also the average of the complex number and its conjugate: $\text{Re}z = (z + \bar{z})/2$. The imaginary part is the displacement projected onto the imaginary axis, or the complex average of the complex number and its reflection across the imaginary axis: $\text{Im}z = (z - \bar{z})/(2i)$.

Polar Form of Complex Numbers

There are physical situations in which a transformation from Cartesian (x, y) coordinates to polar (or cylindrical) coordinates (r, θ) simplifies the algebra that is used to describe the physical problem.

An equivalent coordinate transformation for complex numbers, $z = x + iy$, has an analogous simplifying effect for *multiplicative operations* on complex numbers. It has been demonstrated how the complex conjugate, \bar{z} , is related to a reflection—multiplication is related to a **counter-clockwise** rotation in the complex plane. Counter-clockwise rotation corresponds to increasing θ .

The transformations are:

$$(x, y) \rightarrow (r, \theta) \begin{cases} x = r \cos \theta \\ y = r \sin \theta \end{cases} \quad (8-4)$$

$$(r, \theta) \rightarrow (x, y) \begin{cases} r = \sqrt{x^2 + y^2} \\ \theta = \arctan \frac{y}{x} \end{cases}$$

where $\arctan \in (-\pi, \pi]$.

Multiplication, Division, and Roots in Polar Form

One advantage of the polar complex form is the simplicity of multiplication operations:

DeMoivre's formula:

$$z^n = r^n(\cos n\theta + i \sin n\theta) \quad (8-5)$$

$$\sqrt[n]{z} = \sqrt[n]{z} \left(\cos \frac{\theta + 2k\pi}{n} + i \sin \frac{\theta + 2k\pi}{n} \right) \quad (8-6)$$

Lecture 08 MATHEMATICA® Example 2

Numerical Properties of Operations on Complex Numbers

Download notebooks, pdfs, or html from <http://pruffle.mit.edu/3.016-2006>.

Several examples demonstrate issues that arise when complex numbers are evaluated numerically.

- 1: The relationship $e^{2\pi i} = 1$ is exact.
- 2: However, $e^{2.0\pi i}$ is *numerically* 1.
- 3: Chop removes small evalues that are presumed to be the result of numerical imprecision; it operates on complex numbers as well.
- 5-8 Here, the difference between something that is exactly i and is numerically $1.0i$ is demonstrated...
- 9-12 And, this is similar demostration for $1+i$ using its polar form as a starting point.

```

1 ExactlyOne = Exp[2\pi i]
2 NumericallyOne = Exp[N[2\pi i]]
3 Chop[NumericallyOne]
4 Round[NumericallyOne]
5 ExactlyI = Exp[\pi i/2]
6 NumericallyI = Exp[N[\pi i/2]]
7 Round[NumericallyI]
8 Chop[NumericallyI]
9 ExactlyOnePlusI = ComplexExpand[\sqrt{2} Exp[\pi i/4]]
10 NumericallyOnePlusI = ComplexExpand[\sqrt{2} Exp[N[\pi i/4]]]
11 Chop[NumericallyOnePlusI]
12 Round[NumericallyOnePlusI]
13 Round[1.5 - 3.5 Sqrt[-1]]
14 Re[NumericallyOnePlusI]
15 Im[NumericallyOnePlusI]

```

Exponentiation and Relations to Trigonometric Functions

Exponentiation of a complex number is defined by:

$$e^z = e^{x+iy} = e^x(\cos y + i \sin y) \quad (8-7)$$

Exponentiation of a purely imaginary number advances the angle by rotation:

$$e^{iy} = \cos y + i \sin y \quad (8-8)$$

combining Eq. 8-8 with Eq. 8-7 gives the particularly useful form:

$$z = x + iy = re^{i\theta} \quad (8-9)$$

and the useful relations (obtained simply by considering the complex plane's geometry)

$$e^{2\pi i} = 1 \quad e^{\pi i} = -1 \quad e^{-\pi i} = -1 \quad e^{\frac{\pi}{2}i} = i \quad e^{-\frac{\pi}{2}i} = -i \quad (8-10)$$

Subtraction of powers in Eq. 8-8 and generalization gives known relations for trigonometric functions:

$$\begin{aligned} \cos z &= \frac{e^{iz} + e^{-iz}}{2} & \sin z &= \frac{e^{iz} - e^{-iz}}{2i} \\ \cosh z &= \frac{e^z + e^{-z}}{2} & \sinh z &= \frac{e^z - e^{-z}}{2} \\ \cos z &= \cosh iz & i \sin z &= \sinh iz \\ \cos iz &= \cosh z & \sin iz &= i \sinh z \end{aligned} \quad (8-11)$$

Complex Numbers in Roots to Polynomial Equations

Complex numbers frequently arise when solving for the roots of a polynomial equation. There are many cases in which a model of system's physical behavior depends on whether the roots of a polynomial are real or imaginary, and if the real part is positive. While evaluating the nature of the roots is straightforward conceptually, this often creates difficulties computationally. Frequently, ordered lists of solutions are maintained and the behavior each solution is followed.

Lecture 08 MATHEMATICA® Example 3

Complex Roots of Polynomial Equations

Download notebooks, pdfs, or html from <http://pruffle.mit.edu/3.016-2006>.

Here we construct an artificial example of a model that depends on a single parameter in a quadratic polynomial and illustrate methods to analyze and visualize its roots. Methods to “peek” at the form of long expressions are also demonstrated.

- 1-6 Using a prototype fourth order equation, a list of solutions are obtained; the real and imaginary parts are computed.
- 7: The above is generalized to a single parameter b in the quartic equation; the conditions that the roots are real will be visualized. `bsols`, the list of solution rule-lists is long and complicated.
- 8: First, one must consider the structure of `bsols`, `Dimensions` indicates it is a list of four lists, each of length 1.
- 9: `Short` is a practical method to observe the structure without filling up the screen display.
- 10: `Flatten` converts lists of lists into a single list—it is especially useful with the lists of rule lists that are returned from `Solve`.
- 11: Here, the real and complex parts of *each* of the solutions is obtained with `Re` and `Im` where the parameter b is assumed to be real via the use of `ComplexExpand`. **This may take a long time to evaluate on some computers.**
- 12: Which of the solutions (i.e., 1,2,3, or 4) is identified by using `Hue`.
- 13: Similarly, the real parts appear to converge to a single value when the imaginary parts (from above) appear...
- 14: But, the actual behavior is best illustrated by using `Thickness` to distinguish superposed values. The behavior of real parts of this solution have what is called a *pitchfork structure*.

```

1 sols = Solve[x^4 - x^3 + x + 1 == 0, x]
2 x /. sols
3 Im[x /. sols]
4 ComplexExpand[Im[x /. sols]]
5 ComplexExpand[Im[x /. sols]] // N
6 ComplexExpand[Re[x /. sols]] // N

Generalize the above to a family of solutions: find b such
that imaginary part of the solution vanishes

7 bsols = Solve[(x^4 - x^3 + b*x + 1) == 0, x]
8 Dimensions[bsols]
9 Short[bsols, 4]
10 Dimensions[Flatten[bsols]]
Short[Flatten[bsols], 4]
11 SolsbImag = ComplexExpand[Im[x /. bsols]];
Dimensions[SolsbImag]
Short[SolsbImag[[1]]]
12 SolsbReal = ComplexExpand[Re[x /. bsols]];
13 Plot[Evaluate[SolsbImag], {b, -10, 10}]
14 Plot[Evaluate[SolsbImag], {b, -10, 10},
PlotStyle -> Table[{Hue[1 - a/6]}, {a, 1, 4}]]
15 Plot[Evaluate[SolsbReal], {b, -10, 10},
PlotStyle -> Table[{Hue[1 - a/6]}, {a, 1, 4}]]
16 Plot[Evaluate[SolsbReal], {b, -10, 10},
PlotStyle -> Table[{Hue[1 - a/6], Thickness[0.05 - .01*a]}, {a, 1, 4}]]
17 Plot[Evaluate[SolsbReal], {b, -10, 10}, PlotStyle ->
Table[{Hue[1 - a/6], Thickness[0.05 - .01*a]}, {a, 1, 4}]]
```