

Lecture 26: Separation of Variables and Solutions to Common ODEs

Reading:

Kreyszig Sections: 5.3, 5.5, 5.6 (pages 177–180, 189–197, 198–202)

[3.016 Home](#)



[Full Screen](#)

[Close](#)

[Quit](#)

Special Functions: Solutions to Common ODEs

Most calculators have a button that evaluates the eigensolution to the simple first-order ODE $dy/dt = \lambda y$. Also, most calculators have buttons that evaluate the eigensolutions to the simple second-order ODE: $d^2y/dt^2 = \lambda y$.

Of course, these are also just the exponential and trigonometric functions.

However, there are many more simple differential equations that follow from physical models and these also have known solutions that are not simple combinations of sines, cosines, and exponentials. The solutions to these differential equations are called *special functions*. MATHEMATICA® has an extensive list of special functions and these are collected in its help browser.

For example, the positions of a vibrating drum head are modeled with in cylindrical coordinates by Bessel's equation:

$$\begin{aligned} r^2 \frac{d^2h}{dr^2} + r \frac{dh}{dr} + (k^2 r^2 - m^2)h &= 0 \\ \rho^2 \frac{d^2h}{d\rho^2} + \rho \frac{dh}{d\rho} + (\rho^2 - m^2)h &= 0 \end{aligned} \tag{26-1}$$

where in the second equation $\rho = kr$. The displacement of the drum is $h(r)$; k is related to an inverse wavelength (e.g., the wavelength would be the radius of the drum divided by the number of maxima in the drum head shape) and m is the mode (e.g., the number of maxima traversing the drum by 2π in a circular direction).

There two solutions to Bessel's equation and the general solution is the sum the two:

$$\begin{aligned} h(r) &= C_1 J_m(kr) + C_2 Y_m(kr) \\ h(\rho) &= C_1 J_m(\rho) + C_2 Y_m(\rho) \end{aligned} \quad (26-2)$$

where $J_m(x)$ is called (naturally enough) an *order- m Bessel function of the first kind* and $Y_m(x)$ is called (naturally enough) an *order- m Bessel function of the second kind*. These are analogous to the sines and cosines, but for a different ODE.

Another equation that appears in models of the angular deformations of body in a central force potentials (for example, the ion distribution about a fixed charge; or, the Schrödinger equation for the electron in a hydrogen atom) in spherical coordinates is Legendre's equation:

$$\begin{aligned} \frac{1}{\sin \theta} \frac{d}{d\theta} \left(\sin \theta \frac{d\Xi}{d\theta} \right) + \left[\ell(\ell + 1) - \frac{m^2}{\sin^2 \theta} \right] \Xi &= 0 \\ \frac{d}{d\mu} \left[(1 - \mu^2) \frac{d\Xi}{d\theta} \right] + \left[\ell(\ell + 1) - \frac{m^2}{1 - \mu^2} \right] \Xi &= 0 \end{aligned} \quad (26-3)$$

where $\mu \equiv \cos \theta$ so that $-1 \leq \mu \leq 1$. The value ℓ is related to the number of modes in the θ direction and m is related to the number of modes in the ϕ direction.

Legendre's equation has two solutions:

$$\Xi(\mu) = C_1 P_{lm}(\mu) + C_2 Q_{lm}(\mu) \quad (26-4)$$

The eigensolution $P_{lm}(\mu)$ is called (again, naturally enough) *order m Legendre functions of the first kind* and $Q_{lm}(\mu)$ are called *order lm Legendre functions of the second kind*.

There are many other types of special functions.

[3.016 Home](#)



[Full Screen](#)

[Close](#)

[Quit](#)

Visualizing special functions.

notebook (non-evaluated)

pdf (evaluated)

html (evaluated)

The ODEs that produce Bessel, Legendre, LaGuerre, and Hypogeometric functions are solved and these special functions are visualized.

- 1: This is *Bessel's equation* for $y(x)$.
- 3: *MyPlotStyle* is a function to set *PlotStyle* for a given number of curves so that their colors are spread over *Hue* from red (0) to blue (0.66).
- 4: This will produce at plot of the zeroeth Bessel's function of the first kind (*BesselJ*) and zeroeth Bessel's function of the second kind (*BesselY*)/solutions.
- 6: This is two-parameter form of *Legendre's equation* for $y(x)$.
- 8: This is another form of Legendre's equation for $y(x)$.
- 14: This will produce solutions to *Laguerre's equation* for $y(x)$.

```

Bessel's equation
1 BesselODE = x2 y''[x] + x y'[x] + (x2 - v2) y[x] = 0
2 DSolve[BesselODE, y[x], x]
3 MyPlotStyle[HowMany_Integer] :=
  Table[{Hue[0.66*i/HowMany], Thickness[0.01]}, {i, 0, HowMany}]
4 Plot[{BesselJ[0, x], BesselY[0, x]}, {x, 0, 20}, PlotStyle -> MyPlotStyle[2]]
5 Plot[{BesselJ[1/2, x], BesselY[1/2, x]}, {x, 0, 20}, PlotStyle -> MyPlotStyle[2]]
Legendre's equation
6 LegendreODE = (1 - x2) y''[x] - 2 x y'[x] + (n (n + 1)) y[x] = 0
7 DSolve[LegendreODE, y[x], x]
8 AnotherFormLegendreODE =
  (1 - x2) y''[x] - 2 x y'[x] + (n (n + 1) -  $\frac{m^2}{1 - x^2}$ ) y[x] = 0
9 DSolve[AnotherFormLegendreODE, y[x], x]
10 Plot[{LegendreP[0, x], LegendreQ[0, x]}, {x, -1, 1}, PlotStyle -> MyPlotStyle[2]]
11 Plot[{LegendreP[1, x], LegendreQ[1, x]}, {x, -1, 1}, PlotStyle -> MyPlotStyle[2]]
12 Plot[Evaluate[Table[LegendreP[i, x], {i, 0, 10}]], {x, -1, 1}, PlotStyle -> MyPlotStyle[11]]
13 Plot[Evaluate[Table[LegendreQ[i, x], {i, 0, 10}]], {x, -1, 1}, PlotStyle -> MyPlotStyle[11]]
Hypergeometric and Laguerre special functions
14 DSolve[x y''[x] + (q + 1 - x) y'[x] + p y[x] = 0, y[x], x]
15 Plot[LaguerreL[4, 1, x], {x, -5, 15}]

```

[3.016 Home](#)



[Full Screen](#)

[Close](#)

[Quit](#)

Partial Differential Equations: Separation of Variables

Many *ordinary* differential equations that arise in practice derive from methods to solve *partial differential equations*.

In other words, the solution to the partial differential equation involving $c(x, y, z, t)$ and its partial derivatives with respect to x , y , z , and t can sometimes be reduced to the solution of several ordinary differential equations.

In practice, most of the partial differential equations that can be solved analytically are solved by the *method of separation of variables*. Separation of variables works by isolating one of the variables onto one side of equality—it is best described by simple example and here the one-dimensional wave-equation is a prototype. The wave-equation (e.g., the time (t)-dependent propagation of a scalar quantity (h) such as height, density, charge, etc. in a single direction x is:

$$\frac{\partial^2 h(x, t)}{\partial t^2} = v^2 \frac{\partial^2 h(x, t)}{\partial x^2} \quad (26-5)$$

[3.016 Home](#)[Full Screen](#)[Close](#)[Quit](#)

where v is the phase-velocity $v = \omega/k$, ω is the angular frequency describing how rapidly the phase of the wave changes as it moves past a fixed position and $k = 2\pi/\lambda$ is the wave-number.

Consider a specific case in which waves are propagating in a guitar string of length L —this will give Dirichlet boundary conditions at the guitar's nut and saddle:

$$h(x = 0, t) = 0 \quad \text{and} \quad h(x = L, t) = 0 \quad (26-6)$$

(two boundary conditions—one for each spatial derivative). The shape of plucked string gives the *initial condition*; for example, this could be modeled with a triangular shape:

$$h(x, t = 0) = \begin{cases} Ax/\ell & 0 < x < \ell \\ A\frac{L-x}{L-\ell} & \ell < x < L \end{cases} \quad (26-7)$$

where the string is plucked at a position $x = \ell$ with a displacement A at time $t = 0$.

The separation of variables method begins with the assumption that *the function can be factored into independent functions of the dependent variables*. For Eq. 26-5, this assumption is written as:

$$h(x, t) = \chi(x)\tau(t) \quad (26-8)$$

If this is inserted into Eq. 26-5, and both sides are divided by $v^2\chi(x)\tau(t)$ then

$$\frac{1}{v^2\tau(t)} \frac{d^2\tau(t)}{dt^2} = \frac{1}{\chi(x)} \frac{d^2\chi(x)}{dx^2} \quad (26-9)$$

Note that both sides of Eq. 26-9 depend on different variables and observe Fig. 26-28.




Figure 26-28: If two functions, $T(t)$ and $X(x)$, depend on different variables and are equal, then they can only be constant $\chi(x) = \tau(t) = \lambda$

Thus, both sides of Eq. 26-9 can be set equal to a *separation constant* λ :

$$\frac{1}{\chi(x)} \frac{d^2\chi}{dx^2} = \lambda \quad \text{or} \quad \frac{d^2\chi}{dx^2} - \lambda\chi = 0 \quad (26-10)$$

and

$$\frac{1}{v^2\tau(t)} \frac{d^2\tau}{dt^2} = \lambda \quad \text{or} \quad \frac{d^2\tau}{dt^2} - \lambda v^2\tau = 0 \quad (26-11)$$

[3.016 Home](#)

◀ ▶ ▷ ▸

[Full Screen](#)

[Close](#)

[Quit](#)

For Eq. 26-10, the boundary conditions (Eq. 26-6) can also be written in terms of $\chi(x)$:

$$\chi(x = 0) = 0 \quad \text{and} \quad \chi(x = L) = 0 \quad (26-12)$$

As advertised, ODEs are generated (Eqs. 26-10 and 26-11) in the process of solving the PDE 26-5. As only Eq. 26-10 has a boundary condition, it is solved first; in general its solution appears as:

$$\chi(x) = \begin{cases} A_+ \exp(\sqrt{\lambda}x) + B_+ \exp(-\sqrt{\lambda}x) & 0.25\text{in} \quad \text{if } \lambda > 0 \\ A_0 x + B_0 & \text{if } \lambda = 0 \\ A_- \cos(\sqrt{-\lambda}x) + B_- \sin(\sqrt{-\lambda}x) & 0.25\text{in} \quad \text{if } \lambda < 0 \end{cases} \quad (26-13)$$

The boundary conditions 26-12 place an initial restriction on the separation constant $\lambda > 0$ and specify that $\chi(x)$ must be a sum of sines and cosines. Furthermore, trying to solve Eqs. 26-12 at $x = 0$ shows that $A_- = 0$; at $x = L$, solutions must coincide with any of the *zeroes of the eigenfunction* $\sin(kx) = \sin(\sqrt{-\lambda}x)$, or

$$\lambda_n = \frac{-n^2\pi^2}{L^2} = -k_n^2 \quad n = 1, 2, 3, \dots \quad (26-14)$$

The λ_n (or equivalently the k_n) become *eigenvalues* of the ODE and generate an infinity of eigenfunctions with independent amplitudes A_n : $\chi_n(x) = A_n \sin(n\pi x/L)$.

This infinity of eigenfunctions are needed to satisfy the initial conditions Eqs. 26-7, but first the solution to the second ODE 26-11 must be obtained for the *the restricted set of eigenvalues* for the separation constant:

$$\frac{d^2\tau}{dt^2} + \frac{n^2\pi^2}{L^2} v^2 \tau = 0 \quad (26-15)$$

so that, in general,

$$\tau_n(t) = \mathcal{E}_n^* \cos\left(\frac{n\pi vt}{L}\right) + \mathcal{O}_n^* \sin\left(\frac{n\pi vt}{L}\right) \quad (26-16)$$

and therefore with Eq. 26-8, the superposition of all solutions is

$$\begin{aligned} h(x, t) &= \sum_{n=1}^{\infty} \tau_n(t) \chi_n(x) \\ &= \sum_{n=1}^{\infty} \left(\mathcal{E}_n \cos \frac{n\pi vt}{L} + \mathcal{O}_n \sin \frac{n\pi vt}{L} \right) \sin \frac{n\pi x}{L} \end{aligned} \quad (26-17)$$

[3.016 Home](#)



[Full Screen](#)

[Close](#)

[Quit](#)

And the initial conditions become

$$h(x, 0) = \sum_{n=1}^{\infty} \tau_n(0) \chi_n(x) = \sum_{n=1}^{\infty} \mathcal{E}_n \sin \frac{n\pi x}{L} \quad (26-18)$$

(the sine (odd) coefficients are not needed in this case) and determination of the coefficients is reduced to the Fourier representation. For the initial conditions in Eq. 26-7, these can be computed using Eq. ??:

$$\mathcal{E}_n = \frac{AL^2 \sin \frac{2\pi n \ell}{L}}{(2\pi n)^2 \ell (L - \ell)} \quad (26-19)$$

giving a solution:

$$h(x, t) = \sum_{n=1}^{\infty} \frac{AL^2 \sin \frac{2\pi n \ell}{L}}{(2\pi n)^2 \ell (L - \ell)} \cos \frac{n\pi v t}{L} \sin \frac{n\pi x}{L} \quad (26-20)$$

The Shrödinger equation for a central potential serves as a more involved example for the method of separation of variables and is provided in the following section.

[3.016 Home](#)



[Full Screen](#)

[Close](#)

[Quit](#)

Special Functions in the Eigenfunctions of the Hydrogen Atom

The time-independent Shrödinger for the electron in a hydrogen atom is a partial differential equation involving three spatial variables. If the mass of the nucleus can be considered very large compared to that of an electron, then it is reasonable to fix the center of a spherical $1/r$ -potential at the origin and use spherical coordinates (r, θ, ϕ) to express the Shrödinger equation:¹⁸

$$\frac{\hbar^2}{2m_e} \nabla^2 \psi + V \psi = E \psi \quad (26-21)$$

$$\frac{\hbar^2}{2m_e} \left[\frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial}{\partial r} \right) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2}{\partial \phi^2} \right] \psi - \frac{Ze^2 \psi}{\epsilon_0 r} = E \psi$$

Here, ϕ wraps around like longitude and ϕ goes north and south from the equator ($\theta=0$) like latitude.

¹⁸To treat the hydrogen atom more accurately, the reduced mass $\tilde{m} = 1/(1/m + 1/M)$, weighted coordinates $\tilde{x} = (mx_m + Mx_M)/M$ etc., and relative positions $\Delta x = x_m - x_M$, etc., would produce PDEs for the entire system and for the relative positions.

Because the potential depends only on r , the initial separation is between the radial and angular parts,

$$\psi(r, \theta, \phi) = \rho(r)Y(\theta, \phi) \quad (26-22)$$

which separates Eq. 26-21 into

$$\begin{aligned} \frac{1}{\rho} \frac{d}{dr} \left(r^2 \frac{d\rho}{dr} \right) + \frac{2m_e r^2}{\hbar^2} \left(E + \frac{Ze^2}{\epsilon_o r} \right) \\ = \frac{1}{Y} \left[\frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial Y}{\partial \theta} \right) + \frac{1}{\sin^2 \theta} \frac{\partial^2 Y}{\partial \phi^2} \right] \end{aligned} \quad (26-23)$$

Therefore, the two sides must be equal to a separation constant λ . The radial part becomes

$$\frac{1}{r^2} \frac{d}{dr} \left(r^2 \frac{d\rho}{dr} \right) + \left[\frac{2m_e}{\hbar^2} \left(E - \frac{Ze^2}{\epsilon_o r} \right) - \frac{\lambda}{r^2} \right] \rho = 0 \quad (26-24)$$

and the angular part becomes

$$\frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial Y}{\partial \theta} \right) + \frac{1}{\sin^2 \theta} \frac{\partial^2 Y}{\partial \phi^2} + \lambda Y = 0 \quad (26-25)$$

Solutions to Eq. 26-25 are related to the *spherical harmonics* and can be derived through another separation of variables. Putting

$$Y(\theta, \phi) = \Theta(\theta)\Phi(\phi) \quad (26-26)$$

into Eq. 26-25 gives

$$\frac{1}{\Phi} \frac{d^2 \Phi}{d\phi^2} = -m^2 = \frac{-\sin \theta}{\Theta} \frac{d}{d\theta} \left(\sin \theta \frac{d\Theta}{d\theta} \right) - \lambda \sin^2 \theta \quad (26-27)$$

where the separation constant, $-m^2$, is explicitly set to a negative quantity reflecting that Φ must have periodic solutions, i.e., the two separated ODEs are:

$$\begin{aligned} 0 &= \frac{d^2 \Phi}{d\phi^2} + m^2 \Phi \\ 0 &= \frac{1}{\sin \theta} \frac{d}{d\theta} \left(\sin \theta \frac{d\Theta}{d\theta} \right) \left(\lambda - \frac{m^2}{\sin^2 \theta} \right) \Theta \end{aligned} \quad (26-28)$$

[3.016 Home](#)



[Full Screen](#)

[Close](#)

[Quit](#)

The first of these has the same form as Eq. 26-10 (solutions given by Eq. 26-13) but here the solutions will be written as

$$\Phi(\phi) = A_+ e^{im\phi} + A_- e^{-im\phi} \quad (26-29)$$

because the wavefunction is complex in general. Here, m , indicated how many maxima that the latitudinal part of ϕ will have, and the two different A multiply wavefunctions that are out-of-phase by π . Either solution can be obtained by changing the sign of m , therefore in general

$$\Phi_m(\phi) = \frac{1}{\sqrt{2\pi}} e^{im\phi} \quad m = 0, \pm 1, \pm 2, \dots \quad (26-30)$$

where the prefactor normalizes Φ so that $\int_0^{2\pi} \Phi_m \Phi_m^* d\phi = 1$.

The second of Eqs. 26-28 has the same form as Eq. 26-3 for which the relevant solution (because they are bounded) is an only if

$$\lambda = \ell(\ell + 1) \quad \text{and} \quad |m| \leq \ell \quad \text{for} \quad \ell = 0, 1, 2, \dots \quad (26-31)$$

Putting all of this together and normalizing, the spherical harmonic part of the H-atom orbital (Eq. 26-25) is:

$$Y_{\ell,m} = \sigma(m) \sqrt{\frac{2\ell + 1}{4\pi} \frac{(\ell - |m|)!}{(\ell + |m|)!}} P_{\ell,m}(\cos \theta) e^{im\phi} \quad (26-32)$$

where $\sigma(m) = 1$ if $m \leq 0$ or m even, and $\sigma(m) = -1$ for odd-positive m .

Finally, for the radial part of the H-atom orbital (Eq. 26-24) with $\lambda = \ell(\ell + 1)$ becomes an ODE in r :

$$\frac{1}{r^2} \frac{d}{dr} \left(r^2 \frac{d\rho}{dr} \right) + \left[\frac{2m_e}{\hbar^2} \left(E - \frac{Ze^2}{\epsilon_0 r} \right) - \frac{\ell(\ell + 1)}{r^2} \right] \rho = 0 \quad (26-33)$$

where $E < 0$ defines a bound state.

This equation can also be solved analytically, and it has integer eigenvalues $n > \ell \geq |m|$:

$$\rho_{n,\ell}(r) = -\sqrt{\left(\frac{2Z}{na_0}\right)^3 \frac{(n - \ell - 1)!}{2n[(n + \ell)!]^3}} \exp\left(\frac{-Zr}{na_0}\right) \left(\frac{-2Zr}{na_0}\right)^\ell L_{n+1,2\ell+1}\left(\frac{2Zr}{na_0}\right) \quad (26-34)$$

where $a_0 = \hbar^2 \epsilon_0 / m_e e^2$ is the *Bohr radius* and the $L_{q,p}$ is yet another special function—c'est *LaGuerre polynomials*.

The Hydrogen orbitals are visualized in the following example.

Lecture 26 MATHEMATICA® Example 2

Visualizing the Hydrogen atom eigenfunctions

notebook (non-evaluated)

pdf (evaluated)

html (evaluated)

This example is still in progress—it has not been check for accuracy yet.

1: This example will be completed at a later date.

Functions to display ψ ψ^* will be developed.
Lengths will be normalized so that the Bohr radius is 1 and $Z = 1$.

```

1 x[n_] := x[n] = 1/n
2 A[n_, L_] := A[n, L] = Sqrt[Factorial[n - L - 1]/(2 n (Factorial[n + L])^3)]
3 prefactor[n_, l_] := prefactor[n, l] = A[n, l] (2 x[n])^(3/2)
4 spherefactor[l_, m_] := (2 l + 1 Factorial[l - m])/(4 \[Pi] Factorial[l + m])
5 SquareRadialPart[n_, l_, \rho_] := (prefactor[n, l] Exp[-\rho/2] \rho^l) LaguerreL[n - l - 1, 2 l + 1, 2 \rho]^2
6 SquareAngularPart[l_, m_, prob_, \theta_, \phi_] :=
  Module[{ctheta = Cos[\theta], r, \rho = prob * (spherefactor[l, m] LegendreP[l, m, ctheta])^2, \phi = \phi \[Degree]},
  r = \rho prob * (spherefactor[l, m] LegendreP[l, m, ctheta])^2;
  Return[{r Cos[\phi] ctheta, r Sin[\phi] ctheta, r Sin[\theta]}]
]
7 << Graphics`FilledPlot
8 << Graphics`ParametricPlot3D

```

3.016 Home



Full Screen

Close

Quit

Index

Bessel functions, 220
Bessel's equation, 219, 221
`BesselJ`, 221
`BesselY`, 221
Bohr radius, 228
calculator buttons, 219
 on fancy calculators, 219

drum head, 219

eigenfunctions and eigenvalues, 224

Example function

`MyPlotStyle`, 221

`Hue`, 221

hydrogen atom

 visualizing electron orbitals, 225

initial condition, 222

LaGuerre polynomials, 228

Laguerre's equation, 221

Ledendre's equation, 221

Legendre equation, 220

Legendre functions, 220

Mathematica function

`BesselJ`, 221

`BesselY`, 221

`Hue`, 221

`PlotStyle`, 221
method of separation of variables, 222
`MyPlotStyle`, 221

partial differential equations, 222
phase velocity, 222
`PlotStyle`, 221

separation constant, 223
separation of variables, 222, 225
Shrödinger equation

 central potential, 225

 hydrogen atom, 225

special functions, 219

spherical harmonics, 226

wave equation

 1D, 222

zeroes of the eigenfunction, 224

[3.016 Home](#)



[Full Screen](#)

[Close](#)

[Quit](#)